

AD-A083 698

CARNEGIE-MELLON UNIV PITTSBURGH PA MANAGEMENT SCIENC--ETC F/G 12/1
AN EXTENSION OF SAIGAL'S CLASS OF Q-MATRICES.(U)

FEB 80 J PANG

N00014-75-C-0621


UNCLASSIFIED

MSRR-454

NL

1 OF 1
AD
A083698






①

LEVEL II

AD A083698




Carnegie-Mellon University

PITTSBURGH, PENNSYLVANIA 15213

STIC
S ECTE D
APR 30 1980
E

GRADUATE SCHOOL OF INDUSTRIAL ADMINISTRATION

WILLIAM LARIMER MELLON, FOUNDER



DDC FILE COPY

80 4 30 070

9 Management Science Research Report No. 454

6 AN EXTENSION OF SAIGAL'S CLASS OF Q-MATRICES

by

10 Jong-Shi Pang

14 MSRR-454

11 February, 1980

12 12

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

15
This report was prepared as part of the activities of the Management Sciences Research Group, Carnegie-Mellon University, under Contract N00014-75-C-0621 NR 047-048 with the Office of Naval Research. Reproduction in whole or in part is permitted for any purpose of the U.S. Government.

Management Sciences Research Group
Graduate School of Industrial Administration
Carnegie-Mellon University
Pittsburgh, Pennsylvania 15213

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

403426

mt

A UNIFICATION OF TWO CLASSES OF Q-MATRICES

Jong-Shi Pang

ABSTRACT. This note presents a class of Q-matrices which includes Saigal's class N of Q-matrices with negative principal minors and the class E of strictly semi-monotone Q-matrices.

Key Words: Class of matrices, Linear complementarity problem.

1. Introduction. Given a real square matrix M and a real vector q of the same size, the linear complementarity problem (q, M) is to find a vector x such that

$$q + Mx \geq 0, \quad x \geq 0 \quad \text{and} \quad x^T(q + Mx) = 0.$$

The matrix M is a Q-matrix if the problem (q, M) has a solution for all vectors q .

The problem of constructively identifying a Q-matrix has yet to be solved. Over the past years, numerous classes of matrices have been shown to belong to this large yet very much unknown class of Q-matrices. Two such classes are discovered by Eaves [2] and Saigal [7]. Eaves' class is $L_1 \cap L_2 \cap S$ where L_1 consists of the semi-monotone matrices M which are square matrices such that for each vector $0 \neq x \geq 0$, there is an index k such that $x_k > 0$ and $(Mx)_k \geq 0$; L_2 consists of the square matrices M such that if x is a nonzero solution of the problem $(0, M)$, then there exist nonnegative diagonal matrices D_1 and D_2 with $D_2 x \neq 0$ and $(D_1 M + M^T D_2) x = 0$; and S consists of matrices M for which there is a vector $x > 0$ such that $Mx > 0$. Saigal's class is $N \cap S$ where N consists of square matrices M with negative principal minors. These two classes, namely, $L \cap S$ and $N \cap S$ where $L = L_1 \cap L_2$, are in fact distinct because

$$\begin{bmatrix} -1 & 2 \\ 4 & -1 \end{bmatrix} \in (N \cap S) \setminus L_1 \quad \text{and} \quad \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in (L \cap S) \setminus N.$$

Included among Eaves' class of Q-matrices is the subclass E of strictly semi-monotone matrices which are square matrices M such that for each $0 \neq x \geq 0$, there is an index k with $x_k > 0$ and $(Mx)_k > 0$. The two matrices just given illustrate that the two classes E and $N \cap S$ are not contained in one another.

Our purpose in the note is to present a class of Q-matrices which properly contains the two classes E and $N \cap S$. The construction of this unifying class is very much motivated by the proof used in [7] to establish that $N \cap S \subseteq Q$.

2. Main Results. Let A be an n by n matrix and α an index subset of $\{1, \dots, n\}$. Suppose that the principal submatrix $A_{\alpha\alpha}$ is nonsingular. Let P be a permutation matrix such that $P^T A P$ has $A_{\alpha\alpha}$ as a leading principal submatrix. The α -principal pivot transform of A is defined as the matrix PA^*P^T where

$$(1) \quad A^* = \begin{pmatrix} A_{\alpha\alpha}^{-1} & -A_{\alpha\alpha}^{-1} A_{\alpha\beta} \\ A_{\beta\alpha} A_{\alpha\alpha}^{-1} & A_{\beta\beta} - A_{\beta\alpha} A_{\alpha\alpha}^{-1} A_{\alpha\beta} \end{pmatrix}.$$

Following the notations of Garcia [4], we let $E^*(d)$ for $d > 0$ or $d = 0$, to denote the class of square matrices M for which the problem (d, M) has zero as the unique solution.

We note that A is a Q-matrix if and only if each of its principal pivot transforms is so, i.e., Q-matrices are invariant under principal pivot transforms. Notice however, that the two classes E and $N \cap S$ are not invariant under such transforms.

We say that an n by n matrix A is an \tilde{N} -matrix if there is a vector $f > 0$ and a subset α of $\{1, \dots, n\}$ satisfying the conditions below

- (i) $Af > 0$
- (ii) $A_{\alpha\alpha}$ is nonsingular
- (iii) For any $(n-1)$ by $(n-1)$ principal submatrix \tilde{A} of the α -principal pivot transform of A, it holds that $\tilde{A} \in E^*(\tilde{f}) \cap E^*(0)$ where \tilde{f} is the $(n-1)$ -

subvector of the (positive) vector \bar{f} which in partitioned form (according to A^* given in (1)) is defined as

$$\bar{f} = \begin{pmatrix} f_\alpha \\ A_{\beta\alpha} f_\alpha + A_{\beta\beta} f_\beta \end{pmatrix}.$$

The components of \bar{f} are in correspondence with the rows of \tilde{A}

In the above definition of an \tilde{N} -matrix, we allow $\alpha = \beta$. Of course, the ϕ -principal pivot transform of A is A itself. Note that condition (i) implies that an \tilde{N} -matrix is necessarily an S -matrix. According to [4], the principal submatrix \tilde{A} is a Q -matrix. The result below shows that an \tilde{N} -matrix is in fact a Q -matrix.

Theorem 1. An \tilde{N} -matrix is a Q -matrix.

Proof. Let A be a \tilde{N} -matrix and let A^* be the matrix given in (1). Let q be a given vector and let

$$q^* = \begin{pmatrix} -A_{\alpha\alpha}^{-1} q_\alpha \\ q_\beta - A_{\beta\alpha} A_{\alpha\alpha}^{-1} q_\alpha \end{pmatrix}.$$

It suffices to show that the linear complementarity problem (q^*, A^*) has a solution. We may write this latter problem as

$$x_\alpha = -A_{\alpha\alpha}^{-1} q_\alpha + A_{\alpha\alpha}^{-1} u_\alpha - A_{\alpha\alpha}^{-1} A_{\alpha\beta} x_\beta \geq 0, \quad u_\alpha \geq 0$$

$$u_\beta = q_\beta - A_{\beta\alpha} A_{\alpha\alpha}^{-1} q_\alpha + A_{\beta\alpha} A_{\alpha\alpha}^{-1} u_\alpha + (A_{\beta\beta} - A_{\beta\alpha} A_{\alpha\alpha}^{-1} A_{\alpha\beta}) x_\beta \geq 0, \quad x_\beta \geq 0$$

$$(x_\alpha)^T u_\alpha = (x_\beta)^T u_\beta = 0.$$

Consider the solution of the problem by Lemke's almost complementary

pivoting algorithm [6] using \bar{f} defined in condition (iii) above as the artificial vector. If at some point in the solution process, both x_α and u_β become nonbasic, then the system below has a solution:

$$0 = -A_{\alpha\alpha}^{-1} q_\alpha + \lambda f_\alpha + A_{\alpha\alpha}^{-1} u_\alpha - A_{\alpha\alpha}^{-1} A_{\alpha\beta} x_\beta, \quad u_\alpha \geq 0, \quad \lambda \geq 0, \quad x_\beta \geq 0$$

$$0 = q_\beta - A_{\beta\alpha} A_{\alpha\alpha}^{-1} q_\alpha + \lambda(A_{\beta\alpha} f_\alpha + A_{\beta\beta} f_\beta) + A_{\beta\alpha} A_{\alpha\alpha}^{-1} u_\alpha + (A_{\beta\beta} - A_{\beta\alpha} A_{\alpha\alpha}^{-1} A_{\alpha\beta}) x_\beta.$$

This latter system is clearly equivalent to the one

$$0 = q_\beta + A_{\beta\beta}(x_\beta + \lambda f_\beta), \quad x_\beta \geq 0, \quad \lambda \geq 0$$

$$u_\alpha = q_\alpha - \lambda(A_{\alpha\alpha} f_\alpha + A_{\alpha\beta} f_\beta) + A_{\alpha\beta}(x_\beta + \lambda f_\beta) \geq 0.$$

The consistency of the last system implies that the one below is solvable

$$0 = q_\beta + A_{\beta\beta} \bar{x}_\beta, \quad \bar{x}_\beta \geq 0$$

$$\bar{u}_\alpha = q_\alpha + A_{\alpha\beta} \bar{x}_\beta \geq 0.$$

In fact, we have $\bar{x}_\beta = x_\beta + \lambda f_\beta$ and $\bar{u}_\alpha = u_\alpha + \lambda(A_{\alpha\alpha} f_\alpha + A_{\alpha\beta} f_\beta)$. Therefore in this case, the problem (q, A) has a solution. So suppose that throughout the solution of the problem (q^*, A^*) by Lemke's algorithm (with the above choice of artificial vector \bar{f}), at least one variable in $\begin{pmatrix} x_\alpha \\ u_\beta \end{pmatrix}$ is basic. If the algorithm terminates in a ray, then the problem $(\bar{\lambda}\bar{f}, A^*)$ for some nonnegative $\bar{\lambda}$, has a nonzero solution. This implies by the fact that at least one variable in $\begin{pmatrix} u_\alpha \\ x_\beta \end{pmatrix}$ must be zero, that a certain principal subproblem (\tilde{f}, \tilde{A}) (in the case $\bar{\lambda} > 0$) or $(0, \tilde{A})$ (in the case $\bar{\lambda} = 0$) where \tilde{A} is an $(n-1)$ principal submatrix of A^* and \tilde{f} the corresponding $(n-1)$ -subvector of \bar{f} , would have a nonzero solution. But this contradicts

condition (iii). Consequently, Lemke's algorithm must compute a solution of (q^*, A^*) . This completes the proof of the theorem.

The proof of the theorem is based on an extension of the argument used in Saigal [7] for the special case $N \cap S$. As a matter of fact, the proof also suggests a constructive method for actually computing a solution to (q, A) with A an \tilde{N} -matrix, provided that the index set α and vector f are available readily. Indeed, one can apply Lemke's algorithm to the problem (q^*, A^*) using \bar{f} as the artificial vector. As soon as the artificial variable λ reaches zero or all the u_β and x_α variables become nonbasic, a solution to the given problem (q, A) can be obtained easily.

If a square matrix A is such that some α -principal pivot transform is strictly semi-monotone, then A is an \tilde{N} -matrix. This follows from the fact that principal submatrices of strictly semi-monotone matrices are themselves strictly semi-monotone and that a strictly semi-monotone matrix must be in $E^*(d) \cap E^*(0)$ for any positive d (see [2] e.g.). Hence in particular, if A is itself strictly semi-monotone, then A is an \tilde{N} -matrix.

On the other hand, if a square S -matrix A is such that some α -principal pivot transform \bar{A} has all proper principal minors positive, then A is an \tilde{N} -matrix. This is because any $(n-1)$ by $(n-1)$ principal submatrix of \bar{A} must then be a P -matrix, i.e. has all principal minors positive, and thus belong to $E^*(d) \cap E^*(0)$ for any positive d (see [8]). Hence, in particular, if A is in $N \cap S$ then A is an \tilde{N} -matrix. This follows from the fact that A^{-1} which is the $\{1, \dots, n\}$ -principal pivot transform of A , has all proper principal minors positive (see [7]).

That the class of \tilde{N} -matrices properly contains the union $E \cup (N \cap S)$ can be seen from the example

$$\begin{bmatrix} -1 & 2 \\ -4 & 7 \end{bmatrix}$$

which is an \tilde{N} -matrix (its $\{2\}$ -principal pivot transform is a positive matrix) but certainly not strictly semi-monotone or has negative principal minors.

In [5], it is shown that the linear complementarity problem (q, A) with $A \in N$ has 0, 1, 2 or 3 solutions. The theorem below extends this result. Recall that a square matrix is nondegenerate if all its principal minors are nonzero.

Theorem 2. Let A be a nondegenerate matrix such that some α -principal pivot transform \bar{A} has all proper principal minors positive. Then for every vector q , the linear complementarity problem (q, A) has 0, 1, 2, or 3 solutions.

Proof. If \bar{A} has positive determinant, then \bar{A} and thus A is a P-matrix. Hence the problem (q, A) has a unique solution for all vectors q . On the other hand, if \bar{A} does not have positive determinant, then it must have negative determinant. This is because \bar{A} must be nonsingular. In fact, its inverse is a principal rearrangement of the β -principal pivot transform of A with β the complement of α . This latter principal pivot transform is well-defined by the nondegeneracy of A . Consequently, it follows that the inverse of \bar{A} is in class N . Hence, by the result established in [5], the linear complementarity problem (\bar{q}, \bar{A}^{-1}) has 0, 1, 2 or 3 solutions. As \bar{A}^{-1} is also a principal pivot transform of A , the same conclusion is true for each (q, A) . This completes the proof of the theorem.

We gave an example earlier to show that there are matrices in $N \cap S$ which are not in the class E. The following result establishes that the inverse of a matrix in $N \cap S$ in fact belongs to E.

Theorem 3. Let A be in $N \cap S$. Then the inverse of A is in E.

Proof. In fact, if A is in $N \cap S$, then each proper principal submatrix of A^{-1} is a P-matrix. In particular, each $(n - 1)$ by $(n - 1)$ principal submatrix of A^{-1} is strictly semi-monotone (see [3]). As A^{-1} is also an S-matrix, the desired conclusion now follows from a result established in [1].

REFERENCES

1. R.W. Cottle, "Completely Q-Matrices", Technical Report 79-12, Department of Operations Research, Stanford University (Sept. 1979).
2. B.C. Eaves, "The linear complementarity problem", Management Science 17 (1971) 612-634.
3. D. Gale and H. Nikaido, "The Jacobian matrix and global univalence of mappings", Mathematische Annalen 159 (1965) 81-93.
4. C.B. Garcia, "Some classes of matrices in linear complementarity theory", Mathematical Programming 5 (1973) 299-310.
5. M. Kojima and R. Saigal, "On the number of solutions to a class of linear complementarity problems", Mathematical Programming 17 (1979) 136-139.
6. C.E. Lemke, "Bimatrix equilibrium points and mathematical programming", Management Science 11 (1965) 681-689.
7. R. Saigal, "On the class of complementary cones and Lemke's algorithm", SIAM Journal on Applied Mathematics 23 (1972) 46-60.
8. H. Samelson, R.M. Thrall and O. Wesler, "A partition theorem for Euclidean n-space", Proceedings American Mathematical Society 9 (1958) 805-807.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER M.S.R.R. 454	2. GOVT ACCESSION NO. AD-A083 698	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) AN EXTENSION OF SAIGAL'S CLASS OF Q-MATRICES		5. TYPE OF REPORT & PERIOD COVERED Technical Report February 1980
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Jong-Shi Pang		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0621
9. PERFORMING ORGANIZATION NAME AND ADDRESS Graduate School of Industrial Administration Carnegie-Mellon University Pittsburgh, Pennsylvania 15213		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 047-048
11. CONTROLLING OFFICE NAME AND ADDRESS Personnel and Training Research Programs Office of Naval Research (Code 458) Arlington, Virginia 22217		12. REPORT DATE Februaury 1980
		13. NUMBER OF PAGES 7
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Class of matrices Linear complementarity problem		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This note presents a class of Q-matrices which includes Saigal's class of Q-matrices with negative principal minors and the class of semi-monotone Q-matrices.		

FORM 1 JAN 73 1478

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-66011

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)